

Jacobi polynomials of type BC , Jack polynomials, limit transitions and $O(\infty)$

Tom H. Koornwinder

Extended abstract for the Proceedings of the Conference “Fourier and Radon transformations on symmetric spaces” in honor of Professor S. Helgason’s 65th birthday, Roskilde, Denmark, September 10–12, 1992.

1. Limits of Jacobi polynomials. Let $\alpha, \beta > -1$. Let $p_n^{(\alpha, \beta)}$ ($n = 0, 1, 2, \dots$) be the monic orthogonal polynomials with respect to the measure $(1-x)^\alpha (1+x)^\beta dx$ on the interval $(-1, 1)$ (monic *Jacobi polynomials*). From the explicit expression as hypergeometric series we obtain

$$\lim_{\substack{(\alpha, \beta) \rightarrow \infty \\ \alpha/\beta \rightarrow c}} p_n^{(\alpha, \beta)}(x) = \left(x + \frac{c-1}{c+1} \right)^n.$$

In particular,

$$\lim_{\alpha \rightarrow \infty} p_n^{(\alpha, \alpha)}(x) = x^n. \quad (1)$$

Also,

$$\lim_{\alpha \rightarrow \infty} \alpha^{\frac{1}{2}n} p_n^{(\alpha, \alpha)}(\alpha^{-\frac{1}{2}n} x) = h_n(x), \quad (2)$$

where the h_n are the monic orthogonal polynomials with respect to the measure $e^{-x^2} dx$ on $\mathbb{R}$ (monic *Hermite polynomials*).

2. A representation of $O(\infty)$. Since $p_n^{(\frac{1}{2}d-\frac{3}{2}, \frac{1}{2}d-\frac{3}{2})}$ has an interpretation as spherical function on $O(d)/O(d-1)$, the above limit formulas suggest that monomials and Hermite polynomials have some interpretation on $O(\infty)$. This is indeed the case, see for instance McKean [5] and Matsushima e.a. [4]. Let us explain this briefly. Let S^{d-1} be the sphere of radius $(d/2)^{\frac{1}{2}}$ and midpoint 0 in $\mathbb{R}^d$ and consider it as homogeneous space $O(d)/O(d-1)$. Let π_l^d be the irreducible representation of $O(d)$ on the space $\mathcal{H}_l^d$ of spherical harmonics of degree l on S^{d-1} . For $k < d$ put

$$\mathcal{H}_l^{d,k} := \{f \in \mathbb{C}[x_1, \dots, x_k] \mid f|_{S^{d-1}} \in \mathcal{H}_l^d\},$$

i.e. the space of $O(d-k)$ -invariants in $\mathcal{H}_l^d$ (uniquely) extended to polynomials only depending on the first k coordinates. This space inherits an $O(k)$ -invariant inner product from $\mathcal{H}_l^d$. As $d \rightarrow \infty$, the space $\mathcal{H}_l^{d,k}$ tends to a certain $O(k)$ -invariant inner product space $\mathcal{H}_l^{\infty,k}$ which is a subspace of $\mathbb{C}[x_1, \dots, x_k]$. Let $\mathcal{H}_l^\infty$ be the Hilbert space completion of $\cup_{k=1}^\infty \mathcal{H}_l^{\infty,k}$. Let $O(\infty) := \cup_{d=1}^\infty O(d)$, considered with the inductive limit topology, where, for $d < d'$, $T \in O(d)$ is embedded in $O(d')$ as $\begin{pmatrix} T & 0 \\ 0 & I_{d'-d} \end{pmatrix}$. Then $O(\infty)$ acts on $\mathcal{H}_l^\infty$ by an irreducible unitary representation π_l^∞ .

An orthogonal basis of $\mathcal{H}_l^\infty$ is given by the polynomials

$$\{x \mapsto h_{l_1}(x_1) h_{l_2}(x_2) \dots \mid l_1 + l_2 + \dots = l\}.$$

In particular, if $O_r(\infty)$ denotes the subgroup of all $T \in O(\infty)$ which leave the first r standard basis vectors invariant, then $x \mapsto h_l(x_1)$ is an $O_1(\infty)$ -invariant element of $\mathcal{H}_l^\infty$, unique up to a constant factor. This provides a group theoretic interpretation of (2). The corresponding spherical function can be obtained from the well-known *addition formula* for Hermite polynomials [3, 10.13 (40)]:

$$h_l(x_1 y_1 + \cdots + x_k y_k) = \sum_{l_1 + \cdots + l_k = l} \frac{l!}{l_1! \cdots l_k!} y_1^{l_1} \cdots y_k^{l_k} h_{l_1}(x_1) \cdots h_{l_k}(x_k), \quad y_1^2 + \cdots + y_k^2 = 1.$$

Observe that in the above formula the coefficient of $h_l(x_1)$ on the right hand side equals y_1^l . This yields the spherical function and, moreover, a group theoretic interpretation of (1).

3. A limit for BC_n type Jacobi polynomials. Let R be a root system with Weyl group W and choice of positive roots R^+ , and let $\kappa: R \rightarrow \mathbb{R}$ be a W -invariant mapping. Write $\rho^{(\kappa)} := \frac{1}{2} \sum_{\alpha \in R^+} \kappa(\alpha) \alpha$. Let P^+ be the set of dominant weights, on which a partial order $\leq$ is induced by R^+ . Put $m_\lambda := \sum_{\mu \in W \cdot \lambda} e^\mu$ ($\lambda \in P^+$). The *Jacobi polynomial* associated with R (see Beerends & Opdam [2] and references given there) is an element p_λ ($\lambda \in P^+$) of the form

$$p_\lambda = \sum_{\substack{\mu \in P^+ \\ \mu \leq \lambda}} \Gamma_\mu(\lambda) m_\mu, \quad \Gamma_\lambda(\lambda) = 1,$$

such that it satisfies the differential equation

$$(\text{Laplacian} + \sum_{\alpha \in R^+} \kappa(\alpha) \coth(\alpha/2) \partial_\alpha) p_\lambda = \langle \lambda, \lambda + 2\rho^{(k)} \rangle p_\lambda.$$

For root system BC_n take $R^+ = R_B^+ := \{e_i\} \cup \{2e_i\} \cup \{e_i \pm e_j\}_{i < j}$. Then $P^+ = P_B^+ = \{\lambda \in \mathbb{Z}^n \mid \lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0\}$. Let κ send $e_1, 2e_1, e_1 + e_2$ to k_1, k_2, k_3 , respectively. Write $p_\lambda = p_{B,\lambda}^{(k_1, k_2, k_3)}$.

For root system A_{n-1} take $R^+ = R_A^+ = \{e_i - e_j\}_{1 \leq i < j \leq n}$. Let π be the orthogonal projection which maps $\mathbb{R}^n$ onto the hyperplane $\{t_1 + \cdots + t_n = 0\}$. Then $P_A^+ = \pi(P_B^+)$. Now κ sends R to a fixed $k \in \mathbb{R}$. Write $p_\lambda = p_{A,\lambda}^{(k)}$.

We introduce the *Jack polynomial* here by its expression in terms of A_{n-1} -type Jacobi polynomials:

$$j_\lambda^{(k)}(t) := e^{(t_1 + \cdots + t_n)(\lambda_1 + \cdots + \lambda_n)/n} p_{A,\pi(\lambda)}^{(k)}(\pi(t)), \quad \lambda \in P_B^+, t \in \mathbb{R}^n.$$

The notation used on the left hand side for the Jack polynomial, is not the standard one. The following limit result generalizes (1).

Theorem 1 (Beerends & Koornwinder [1]). Let $-\infty \leq a \leq 1$. Let $\tau, t \in \mathbb{R}^n$ be related by $e^{\tau_i} = (a-1)(a-2)^{-1} + (\sinh(\tau_i/2))^2$. Then

$$\lim_{\substack{(k_1, k_2) \rightarrow \infty \\ k_1/k_2 \rightarrow -a}} p_{B,\lambda}^{(k_1, k_2, k_3)}(t) = \text{const. } j_\lambda^{(k_3)}(\tau).$$

4. An interpretation of Jack polynomials on $O(\infty)$. We first summarize some results of Olshanskiĭ [6]. Denote the left upper $r \times r$ submatrix of the matrix $T \in O(\infty)$ by $\Theta_r(T)$. Then the mapping which sends a double coset $O_r(\infty).T.O_r(\infty)$ to $\Theta_r(T)$ is well-defined and gives a bijection of the double coset space $O(\infty)//O_r(\infty)$ onto the semigroup $\Gamma(r)$ of all $r \times r$ matrices with norm ≤ 1 . Let π be an irreducible unitary representation of $O(\infty)$ on a Hilbert space $\mathcal{H}$ such that the subspace $\mathcal{H}_r$ of $O_r(\infty)$ -fixed elements in $\mathcal{H}$ contains nonzero elements. Let P_r be the orthogonal projection of $\mathcal{H}$ onto $\mathcal{H}_r$.

Theorem 2 (Olshanskiĭ). The mapping $\Theta_r(T) \mapsto P_r\pi(T)|_{\mathcal{H}_r}$ ($T \in O(\infty)$) is a semigroup representation of $\Gamma(r)$ on $\mathcal{H}_r$. Moreover, the space $\mathcal{H}_r$ has finite dimension and the semigroup representation has a unique extension to an irreducible group representation of $GL(n, \mathbb{R})$ on $\mathcal{H}_r$.

Suppose now that the space of $O(r) \times O_r(\infty)$ -invariant elements of $\mathcal{H}$ has dimension one. Then we can speak about the corresponding spherical function on $O(\infty)$ with respect to $O(r) \times O_r(\infty)$. By the above Theorem this spherical function can be immediately connected with a spherical function on $GL(r, \mathbb{R})$ with respect to $O(r)$ corresponding to a finite-dimensional irreducible representation of $GL(r, \mathbb{R})$. These last spherical functions were called *zonal polynomials* by A. T. James and they are special instances of Jack polynomials.

For certain values of k_1, k_2, k_3 the BC_r -type Jacobi polynomial can be interpreted as a spherical function on $O(d)$ with respect to $O(r) \times O(d-r)$. It is plausible to suppose that such a spherical function tends, for $d \rightarrow \infty$, to a certain spherical function on $O(\infty)$ with respect to $O(r) \times O_r(\infty)$. When we connect this, in its turn, to a spherical function on $GL(r, \mathbb{R})$ with respect to $O(r)$ and write this as a Jack polynomial then we would arrive at a group theoretic interpretation of Theorem 1 in special cases.

References

- [1] R. J. Beerends & T. H. Koornwinder, *Analysis on root systems: A_{n-1} as limit case of BC_n* , in preparation.
- [2] R. J. Beerends & E. M. Opdam, *Certain hypergeometric series related to the root system BC* , Trans. Amer. Math. Soc., to appear.
- [3] A. Erdélyi, W. Magnus, F. Oberhettinger & F. G. Tricomi, *Higher transcendental functions, Vol. II*, McGraw-Hill, 1953.
- [4] H. Matsushima, K. Okamoto & T. Sakurai, *On a certain class of irreducible unitary representations of the infinite dimensional rotation group I* , Hiroshima Math. J. 11 (1981), 181–193.
- [5] H. P. McKean, *Geometry of differential space*, Ann. Probab. 1 (1973), 197–206.
- [6] G.I. Olshanskiĭ, *Representations of infinite-dimensional classical groups, limits of enveloping algebras and Yangians*, in: *Topics in representation theory*, A. A. Kirillov (ed.), Advances in Soviet Mathematics 2, 1991.

University of Amsterdam, Faculty of Mathematics and Computer Science
Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands
email `thk@fwi.uva.nl`